



Linking temperate demersal fish species to habitat: scales, patterns and future directions

Andrew F Johnson, Stuart R Jenkins, Jan G Hiddink & Hilmar Hinz

School of Ocean Sciences, Bangor University, Menai Bridge, Anglesey, LL59 5AB, UK

Abstract

Adoption of the ecosystem approach to fisheries management relies on recognition of the link between fish and other components of the ecosystem, namely their physical and biological habitat. However, identifying the habitat requirements of marine fishes and hence determining their distribution in space and time is scientifically complex. We analysed the methodologies and findings of research on temperate, demersal fish habitat requirements to highlight the main developments in this field and to identify potential shortfalls. Many studies were undertaken over large spatial scales (≥ 100 km²) and these generally correlated abundances of fish to abiotic variables. Biological variables were accounted for less often. Small spatial scale (≤ 1 km²), experimental studies were comparatively sparse and commonly focused on biotic variables. Whilst the number of studies focusing on abiotic variables increased with increasing spatial scale, the proportion of studies finding significant relationships between habitat and fish distribution remained constant. This mismatch indicates there is no justification for the tendency to analyse abiotic habitat variables at large spatial scales. Innovative modelling techniques and habitat mapping technologies are developing rapidly, providing new insights at the larger spatial scales. However, there is a clear need for a reduction in study scale, or increase in resolution additional to the integration of biotic variables. We argue that development of sound predictive science in the field of demersal fish habitat determination is reliant on a change in focus along these lines. This is especially important if spatial management strategies, such as Marine Protected Areas (MPA) or No Take Zones (NTZ), are to be used in future ecosystem-based approaches to fisheries management.

Correspondence:

Andrew F Johnson,
School of Ocean
Sciences, Bangor
University, Menai
Bridge, Anglesey
LL59 5AB, UK
Tel.: +44 (0) 1248
388 209
Fax: +44 (0)1248
716 367
E-mail: a.johnson@
bangor.ac.uk; andrew
johnson540@hotmail.
com

Received 14 Sep 2011
Accepted 16 Feb 2012

Keywords abiotic and biotic habitat descriptors, conservation, demersal fish, ecosystem approach, essential fish habitat, habitat association.

Introduction	2
Methods	4
Search criteria	4
Statistical analyses and graphical outputs	4
Results	5
Discussion	10
Data limitations and considerations	10
Spatial scale	11

Abiotic habitat variables	14
Biotic habitat variables	14
Temporal aspects of habitat use	15
Ontogenetic changes in habitat use	15
Non-commercial and rare species	16
Habitat suitability modelling	16
Technological advancements	16
Conclusion	17
Acknowledgements	18
References	18

Introduction

Resource exploitation, habitat modification, pollution and climatic shifts are having widespread and detrimental consequences for marine ecosystems globally (Vitousek 1997; Jackson *et al.* 2001; Kaiser *et al.* 2002). With more than 20% of the world's human population living in biodiversity hotspots (Cincotta *et al.* 2000) and over 60% living within 60 km of the coast (Alongi 1998), marine ecosystems are now under huge pressure from anthropogenic activities and increasingly demonstrate a reduced capacity to withstand these pressures (Folke *et al.* 2004). Of particular concern to managers and conservationists are the impacts of bottom trawl fisheries on benthic ecosystems. The modification and degradation of habitats associated with bottom fishing can have long-term consequences for benthic biota, reducing productivity, biomass and species richness (Hiddink *et al.* 2006; Queiros *et al.* 2006; Hinz *et al.* 2009). Following growing concerns regarding the long-term sustainability of marine fish exploitation (Pauly *et al.* 2005), fisheries collapse from food web restructuring (Frank *et al.* 2005) and reduced abilities of chronically impacted benthic communities to sustain previous stock levels (Shephard *et al.* 2010; Hiddink *et al.* 2011), traditional and often single-species management practices are being evaluated. The result has been the development of a new ecosystem-based fisheries management paradigm (Pikitch *et al.* 2004; Frid *et al.* 2005) which aims to maintain a sustainable ecosystem state by integrating knowledge of biotic, abiotic and human components of ecosystems and applying them to fisheries within ecologically meaningful boundaries (Link 2002).

Major drivers of this new approach have been the formation of numerous conventions and legislations

that have highlighted the importance of defining habitat requirements of key species. Examples include the Conservation of Antarctic Marine Living Resources (1982) and the Sustainable Fisheries Act (SFA) (1996) in the United States. The SFA added Essential Fish Habitat (EFH) provisions to the existing Magnuson–Stevens Fishery Convention (1976) requiring the National Oceanographic and Atmospheric Administration (NOAA) to facilitate the long-term protection of 'those waters and substrates necessary for spawning, breeding, feeding or growth to maturity'. The SFA highlights the importance of defining physical, chemical and biological properties that are used by fish and those that are necessary to support a sustainable fish biomass for each of the fish species listed. Similar initiatives aiming to define the habitat requirements of marine fish species are now being echoed globally [Europe – Directive on the Conservation of Natural Habitats and Wild Fauna and Flora (1992/43/EEC), Australia – Environment Protection and Biodiversity Conservation Act 1999, Canada – Fisheries Act (reforms – sect. 35, 1985) and Species At Risk Act (Bill C-5, 2002)].

Despite the drive towards ecosystem conservation, and long histories of commercial fishing in Europe, Canada and America (Pauly *et al.* 2005), still relatively little is known about the habitat requirements and basic ecology of many temperate demersal fish species (Benaka 1999). With declining marine fish stock levels (Pauly *et al.* 2005) and an increased awareness of the impacts of bottom fisheries (Kaiser 1998), understanding the specific habitat requirements of demersal fishes is more important than ever. Management strategies aiming to attain sustainable exploitation levels of fish stocks can vary widely, from fishing quota implementation (Karagiannakos 1996) or the establishment of no

take zones (NTZ) and marine protected areas (MPA) (Gaines *et al.* 2010), to stock restoration projects involving artificial habitat construction (Seaman 2008) and re-stocking (Heenan *et al.* 2009). The latter initiatives are often only feasible over certain habitat types and scales. For example, artificial habitat construction is a relatively straightforward process for tropical reef fish species, which often have small home ranges and require complex three-dimensional habitats (Roberts and Ormond 1987), and has been implemented successfully in numerous locations over generally small spatial scales (Seaman 2008). Temperate systems on the other hand often consist of relatively homogeneous, soft sediment habitat types that may span thousands of square kilometres over which habitat manipulation is not practicable.

Spatial management measures, such as NTZ's and MPA's, can range from access restrictions for certain users at specified times to full protection from all anthropogenic activity within the designated area. These may vary from closing areas greater than 1000 km² for the protection of multispecies communities with relatively low habitat specificity and wide distributions as illustrated by ground fish stocks in the Western Gulf of Maine, New England, USA (see Murawski *et al.* 2000) to the closures of smaller areas less than 100 km² to deal with single species in areas of high habitat suitability such as the Red mullet (*Mullus barbatus*, Mullidae) in the Gulf of Castellamare, northwestern Sicily, Italy (see Fiorentino *et al.* 2008). Such approaches are now common place; however, their aims with regard to fisheries may vary from the protection of certain life stages of specific species to the overall enhancement of multiple fisheries (Koss *et al.* 2005; Frost and Andersen 2006; Ardron 2008; Gelcich *et al.* 2009). Studies that accurately describe the habitat requirements of target species and the scales over which they operate can facilitate decisions on the type of management strategies that are most viable for the species of concern.

To aid in the sustainable exploitation of demersal fish stocks, researchers must uncover the most important habitat variables determining the distributions of species as well as the range of temporal and spatial scales over which they operate (Hinze *et al.* 2003). Many studies in this field, however, use widely spaced sampling stations (10s–100s km) and combine data collected across temporal scales. Consequently, habitat variables influencing demer-

sal fish habitat distributions that operate over smaller scales may well be overlooked. For example, combinations of monthly to annual data are likely to mask differences in habitat use between seasons. Similarly, collection of data over a scale of hundreds of square kilometres may not allow for inference about differences in distributions at smaller spatial scales. Mismatches between spatial scales examined and the actual scale at which processes occur may at best mean that patterns are missed, and at worst result in drawing erroneous conclusions. Management interventions supported by such outputs may therefore produce questionable management strategies and a lack of sustainable and efficient protection.

The determination of EFH commonly involves relating the abundance of the species under investigation to measured habitat variables thought to be important to the species distribution (Rice 2005). Habitats accommodating the study species are assumed to be of some importance. Stoner (2003), however, points out that seemingly appropriate habitats may never be occupied, with specific locations being more important than particular habitat forms. Ideally, the determination of EFH should be based on more than abundance alone and should also address the growth and survival of individuals, thus considering how their current distribution may affect future generations. This is especially important if predictions derived from research are to be used in management. Such approaches combining measures of growth and fitness are, however, often restricted to small-scale manipulative experiments, as such data collection over large areas is often constrained by methodologies and cost.

Although work in the field of demersal fish habitat determination is becoming an important part of the ecosystem-based approach to fisheries management, we are currently lacking a critical synthesis that assesses the importance of biotic and abiotic habitat variables at different spatial scales. This means that we are currently unaware whether there are particular spatial scales over which different habitat variables affect the distribution of demersal fishes. In this paper, we analyse the number of different habitat variables that have been studied and the spatial scales over which they determine the distribution of temperate, demersal fishes from square metres to more than a hundred thousand square kilometres. We restrict our analysis to temperate, demersal species as the habitat

variables determining the distribution of tropical and pelagic species are likely to differ considerably (see Shepherd and Litvak 2004). Over the period 1980–2011, we aim to quantify the following parameters in this field of research: (i) the number of studies in the field of demersal fish habitat identification and the geographic locations in which they were undertaken, (ii) the life stages and fish species analysed, (iii) the types and numbers of habitat variables commonly investigated and the spatial scale of work undertaken, (iv) the proportion of studies reporting significance for each habitat variable over the spatial range studied and (v) the temporal scales considered in the literature. To examine whether trends in research were justified, we compared the relationships between the total number of studies and the proportion of studies reporting significance for each separate habitat variable, as well as for all abiotic and all biotic variables, across the range of spatial scales analysed. Our analyses permit a critical evaluation of current trends and patterns in the field of demersal fish habitat determination. With rising expectations of managers to meet sustainability targets and the growing pressures on marine ecosystems, continued research in the field of demersal fish habitat determination is required if we are to sustainably exploit our bottom fisheries resources. This analysis is therefore both timely and necessary to assess what research has been undertaken in this field to date, in order that we drive future research efforts in the appropriate direction.

Methods

Search criteria

We conducted a literature search for peer-reviewed publications, published in English over the period 1980–2011, describing the habitat requirements of temperate, marine, demersal fish species. Within any particular publication, one or a number of different investigations may be described. We therefore define ‘article’ as a published journal paper and ‘study’ refers to the separate investigations of different habitat variables undertaken within an article. The period was chosen due to the accessibility of online articles and because it corresponded approximately with the time over which fisheries science wholly embraced research into the anthropogenic impacts on demersal resources and demersal fish habitat determination [following the

announcement of the Magnuson–Stevens Fishery Convention (1976)]. The literature search was undertaken using the commercial search engine Google Scholar, which indexes the full text of scholarly literature across an array of publishing formats. Combinations of the following search terms were used: ‘association’, ‘choice’, ‘connectivity’, ‘demersal’, ‘determining’, ‘distribution’, ‘ecology’, ‘EFH’, ‘essential’, ‘fish’, ‘habitat’, ‘requirement’, ‘scale’, ‘suitability’ and ‘use’. Google Scholar was chosen over Web of Science and Scopus as it gives a higher number of results per general search term (on average), covers non-ISI listed journals (wider search base), and gives lower citation noise [lower citation variation (85% unique entries compared with ISI’s 60%)] (Belew 2005; Pauly and Stergiou 2005; Meho and Yang 2007; Harzing and Van der Wal 2008).

The first one hundred search results from each keyword combination were examined. From these, only peer-reviewed articles were chosen as they form the main body of literature widely available to researchers. The articles used therefore did not include any university theses, technical governmental reports or conference proceedings. Articles examining larval life stages were excluded from the analysis as they form a significant and separate body of literature. Often these early life stages also show markedly different habitat preferences to post-larval developmental stages (Gibson 1994; Sullivan *et al.* 2006). Articles investigating survivorship, condition or fitness of demersal species under differing habitat conditions were included in the database when they were directly related to habitat preferences of the species. From each article fitting the above search criteria, eight categorical data variables were extracted and entered into a database (see Table 1 for details). The variables were publication year, location and total spatial scale of study, life stages, fish species, habitat variables and time trends analysed, and origin of data utilized. The total spatial scale (total area) of each study was calculated using ARC GIS (version 9.2) if it was not described clearly by the authors within the methods. If replicate sites were tested, the area of each was calculated and the total across all sites used. Data were categorized by their origin into stock assessment, field study and laboratory study to illustrate the origin of data sources within this field of research. Habitat variables investigated by authors were classified into abiotic vs. biotic and further into the sub-categories listed in Table 1.

Table 1 Table summarizing the data extracted from each article fitting the search criteria and entered into the first database.

Parameter	Description of parameter	Database input categories
Year	Year of publication	1980–2010 (1 year intervals)
Location	Geographic location (within temperate zones) of work undertaken	N. America, Europe, Australia, New Zealand, S. America, other
Total study scale	Total area of the study site (both laboratory and field) over which replicates were taken	$\leq 1e^{-6} \text{ km}^2$ (cm^2), $\leq 1 \text{ km}^2$ (m^2), $\leq 10 \text{ km}^2$, $\leq 100 \text{ km}^2$, $\leq 1000 \text{ km}^2$, $\leq 10\,000 \text{ km}^2$, $\leq 100\,000 \text{ km}^2$
Life stages analysed	Ontogenetic stage considered in the analyses	Adult, Juvenile, All
Fish order analysed	Fish species and order under study	Mixed*, Gadiformes, Pleuronectiformes, Perciformes, Scorpaeniformes, Carcharhiniformes
Study type	Origin of data used	Laboratory study, Field study, Stock assessment survey
Habitat variables analysed	Variables analysed/hypothesized to be important to the distribution/habitat requirements	Abiotic: depth, hydrography, other abiotic ^a , salinity, substrate, temperature Biotic: competition ^b , biogenic complexity ^{c,†} , predation ^{d,†}
Time trends analysed	Temporal trends formally analysed as part of the hypotheses	Not applicable, day (12 h), day vs. night (24 h), month (30 days), season (90 days), annual (365 days)

*Mixed species are defined as studies looking at four or more different demersal fish species; ^aIncludes dissolved oxygen/level of hypoxia and light; ^bIncludes intra and interspecific competition; ^c3D habitat structures of biological origin; ^dPredation of the demersal species under investigation. [†]Often exposure to predation is a result of biogenic complexity. The distinction between the two was made by referring to the authors' hypotheses and final conclusions.

Statistical analyses and graphical outputs

Initially, we focused on the distribution of journal articles across the years and spatial scales they addressed. Ordinary least squares (OLS) linear regressions were employed to analyse the relationships between (i) the number of peer-reviewed journal articles published in the field and the year of publication, (ii) the number of studies and total spatial scale of each study.

Next, we examined the relationship between the number of studies conducted at particular scales and the proportion of studies that reported significant results to understand whether the concentration of studies at particular scales was justified. Here, only articles which reported results based on individual species were used because those which only reported relationships for mixed assemblages could potentially mask significant relationships at the species level. Regressions of spatial scale vs. number of studies were undertaken for all of the different habitat variables plus 'Total abiotic' and 'Total biotic' categories. This process excluded geography, hydrography and predation as there were too few studies that investigated these variables. This was then repeated, except the number of

studies was replaced with the proportion of these studies which showed a significant relationship.

Data were \log_{10} transformed to ensure normal distributions where required. To test whether there was a significant difference between the spatial scales at which abiotic and biotic habitat variables operate, Wilcoxon rank sum *t*-tests were used.

Results

The literature search identified 109 peer-reviewed journal articles determining the habitat requirements of demersal fish species from the period between 1980 and 2011 (Table 2). The number of articles published increased significantly over time ($R^2 = 0.506$, $F_{1,29} = 29.72$, $P \leq 0.001$); 43% were published since 2005, with the highest number of articles published in 2001 (13), followed by 2009 (12) and 2002/2007 (11 each) (Fig. 1). North America (46%) and Europe (38%) dominated the locations where studies had been undertaken. The remaining 16% of studies were carried out in Australia, New Zealand and South America. No publications came from Africa or Asia.

The majority of studies (67%) made no clear distinction between adult and juvenile life stages

Table 2 Table summarizing separate fish species, life stages and habitat variables analysed.

Fish order	Fish species	Life stage analysed		Habitat variables analysed											Reference		
		Adult	Juvenile	Abiotic						Biotic							
				Depth	Geography	Hydrography	Salinity	Substrate	Temperature	Other	Competition	Biogenic complexity	Predation	Prey resource			
Carcharhiniformes Gadiformes	<i>Scyliorhinus canicula</i>	*	*								*	*	*	*	*		Sims et al. (2001)
	<i>Gadus morhua</i>				*	*					*	*	*	*	*	*	Borg et al. (1997), Bjornstad et al. (1999), Fromentin et al. (2001), Lindholm et al. (2001), Blanchard et al. (2005), Robichaud and Rose (2006)
Perciformes	<i>Gadus macrocephalus</i>		*								*						Laurel et al. (2007, 2009)
	<i>Eleginus gracilis</i>		*								*						Laurel et al. (2007, 2009)
	<i>Merluccius bilinearis</i>	*	*			*					*						Auster et al. (2003)
	<i>Merluccius merluccius</i>		*				*				*						Cartes et al. (2009)
	<i>Cheilodactylus spectabilis</i>		*														McCormick (1998)
	<i>Micropogonius undulatus</i>		*														Eby et al. (2005)
	<i>Pagrus auratus</i>		*														Thrush et al. (2002)
	<i>Paralabrax clathratus</i>	*	*									*					Anderson (2001)
	<i>Sciaenops ocellatus</i>		*									*					Bachelet et al. (2009)
	<i>Tautoglabrus adspersus</i>		*												*		Levin (1994)
Pleuronectiformes	<i>Buglossidium luteum</i>		*			*											Amara et al. (2004)
	<i>Hippoglossoides platessoides</i>		*			*									*		Walsh et al. (2004)
	<i>Limanda ferruginea</i>		*			*								*			Simpson and Walsh (2004), Walsh et al. (2004)
			*			*								*			

Table 2 Table summarizing separate fish species, life stages and habitat variables analysed.

Fish order	Fish species	Life stage analysed	Habitat variables analysed													Reference	
			Abiotic						Biotic						Prey resource		
			All	Juvenile	Adult	Depth	Geography	Hydrography	Salinity	Substrate	Temperature	Other	Competition	Biogenic complexity			Predation
	<i>Hippoglossoides stenolepis</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	Stoner and Abookire (2002), Stoner and Titgen (2003), Ryer et al. (2004), 2007
	<i>Lepidopsetta polyxystra</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	Stoner and Titgen (2003), Ryer et al. (2004), Stoner et al. (2007)
	<i>Limanda limanda</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	Gibson and Ezzi (1987), Hinz et al. (2005)
	<i>Pleuronectes americanus</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	Manderson et al. (2000), (2002), (2003), (2006), Phelan et al. (2001), Stoner et al. (2001)
	<i>Pleuronectes platessa</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	Gibson and Robb (1992), Le Pape et al. (2003), Hinz et al. (2006), Shucksmith et al. (2006), Maxwell et al. (2009)
	<i>Solea solea</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	Eastwood et al. (2003), Le Pape et al. (2003), Hinz et al. (2006), Vinagre et al. (2006), Nicolas et al. (2007), Maxwell et al. (2009)
	<i>Microstomus kitt</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	Hinz et al. (2006), Vinagre et al. (2006)
	<i>Solea senegalensis</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	Vinagre et al. (2006)
Scorpaenif-ormes	<i>Sebastes alutus</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	Brodeur (2001)
	<i>Sebastes</i> sp.	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	Johnson et al. (2003)
Rajiformes	<i>Raja clavata</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	Maxwell et al. (2009)

Table 2 Table summarizing separate fish species, life stages and habitat variables analysed.

Fish order	Fish species	Adult	Juvenile	All	Habitat variables analysed											Reference
					Abiotic					Biotic						
					Depth	Geography	Hydrography	Salinity	Substrate	Temperature	Other	Competition	Biogenic complexity	Predation	Prey resource	
Mixed	<i>Mixed</i>	*	*	*	*	*	*	*	*	*	*	*	*	*	*	Iglesias (1981), Macpherson (1981), Sedberry and Vandolah (1984), Mahon and Smith (1989), De Ben et al. (1990), Ekau (1990), Pihl et al. (1991), Gabriel (1992), Gibson (1994), Koslow et al. (1994), Edgar and Shaw (1995), Fujita et al. (1995), Syms (1995), Gutt and Ekau (1996), Farina et al. (1997), McClatchie et al. (1997), Moranta et al. (1998), Piet et al. (1998), Gaertner et al. (1999), Hyndes et al. (1999), Abookire et al. (2000), Dean et al. (2000), Demestre et al. (2000), Ellis et al. (2000), McConnaughey and Smith (2000), Amezcua and Nash (2001), Auster et al. (2001), Hindell et al. (2001), Methven et al. (2001), Williams and Bax (2001), Araujo et al. (2002), Blagi et al. (2002), Francis et al. (2002), Hughes et al. (2002), Lazzari and Tupper (2002), Magnussen (2002), Kaiser et al. (2004), Pittman et al. (2004), Busby et al. (2005), Logerwell et al. (2005), Massuti and Renones (2005), Perry et al. (2005), Leathwick et al. (2006), Sullivan et al. (2006), Wilhelmsson et al. (2006), Anderson and Yoklavich (2007), Medina et al. (2007), Stal et al. (2007), Bergstad et al. (2008), Selleslagh and Amara (2008), Anderson et al. (2009), Ehrich et al. (2009), Katsanevakis and Maravelias (2009), Katsanevakis et al. (2009), Laidig et al. (2009), Langhamer and Wilhelmsson (2009), Moore et al. (2009), (2011), Ordines and Massut (2009), Persohn et al. (2009), Quiroga et al. (2009), Chatfield et al. (2010), Colloca et al. (2010), Damalas et al. (2010), Keller et al. (2010), Reiss et al. (2010)




Figure 1 Number of peer-reviewed journal articles that have investigated the habitat requirements of temperate demersal fishes (by year of publication). Studies are separated into the major life stages analysed.

analysed. Juvenile life stages were analysed in 26% of articles whilst 7% studied only the adult life stage (Fig. 1). Mixed-species communities were studied in 57% of the studies analysed. Of the remaining 42%, Pleuronectiformes dominated the literature (52%) followed by Gadiformes (25%) and Perciformes (15%) (Table 2). The most studied single species were the Atlantic cod (*Gadus morhua*, Gadidae), Common sole (*Solea solea*, Soleidae) and Winter flounder (*Pseudopleuronectes americanus*, Pleuronectidae). Ninety per cent of all the non-mixed species studied are currently commercially exploited.

Few studies determining the habitat requirements of temperate demersal fishes came from laboratory studies (8%) with the majority originating from field and stock assessment-derived data (46% each respectively) (Fig. 2). Stock assessment-derived data were predominantly (98%) focussed over medium to large spatial scales of study (100s–100 000s km²), field study-derived data (85%) over medium scales (10s–100s km²) and laboratory scale data (100%) over the two smallest scales (cm² to m²).

We found that the majority of studies focussed on the effects of abiotic (72%) rather than biotic (28%)

habitat variables (Fig. 3). There was a tendency for abiotic variables to be studied at larger spatial scales ($R^2 = 0.753$, $F_{1,6} = 23.160$, $P = 0.003$). However, no significant relationship between the number of studies analysing biotic habitat variables and the spatial scales of study was demonstrated ($R^2 = 0.046$, $F_{1,6} = 0.305$, $P = 0.6$). Overall, there was a significant difference between the scales at which abiotic and biotic habitat variables were studied ($P < 0.001$, $W_s = 5051$); abiotic variables focussed at larger spatial scales (69% at scales ≥ 100 km²) and biotic variables at low to medium spatial scales (53% at scales ≤ 100 km²) (Fig. 3).

Assessment of how the proportion of studies showing significant relationships changed with spatial scale was conducted for each habitat variable separately and for the categories 'Total abiotic' and 'Total biotic' (Figs 4 and 5). For the abiotic variables, there was a significant positive relationship between the spatial scale of study and the total number of studies for depth ($R^2 = 0.845$, $F_{1,5} = 35.207$, $P = 0.001$), temperature ($R^2 = 0.767$, $F_{1,4} = 19.766$, $P = 0.004$) and for 'Total abiotic' ($R^2 = 0.846$, $F_{1,7} = 32.990$, $P = 0.001$) (Fig. 4). However when each of these variables and variable combinations was analysed using the




Figure 2 Number of studies originating from laboratory studies, field studies and stock assessments by total spatial scale of study.

proportion of studies which showed a significant relationship, rather than total number of studies, no significant regressions were found. Thus, whilst the total number of studies increased with spatial scale, there was no apparent increase in the proportion of studies showing significant relationships (Fig. 4). For the 'Total biotic' category, there was no clear pattern of change in the number of studies with scale, although one biotic habitat variable (competition) showed an increase in the number of studies with increasing spatial scale ($R^2 = 0.663$, $F_{1,5} = 11.812$, $P = 0.014$) (Fig. 5). There were no significant relationships between the proportion of significant studies and spatial scales of study for any of the individual biotic habitat variables or 'Total biotic' category.

Examination of the distribution of studies among the different habitat variables showed that depth and substrate made up more than half of the abiotic variables analysed (33 and 26%, respectively) (Fig. 4). The most studied biotic habitat variable was biogenic complexity (48%) (Fig. 5). The variables of geography and hydrography, which formed 7 and 0.9% of abiotic habitat variables, respectively,

and competition and predation, which made up 7 and 5% of biotic habitat variables, respectively, were seldom analysed.

Temporal trends were formally analysed in 46% of studies (Fig. 6). Annual and monthly differences in habitat requirements were the most common temporal trends analysed in studies (13% each), followed by season (11%) and day (9%).

Discussion

Data limitations and considerations

Our comprehensive literature search has allowed us to highlight some important trends and patterns in the identification of demersal fish habitat. It is, however, important to discuss possible limitations of our approach and the bearing they may have on our conclusions. This quantitative, systematic analysis only included peer-reviewed articles. The inclusion of dissertations, theses, government reports and conference proceedings may have added to the total number of articles reviewed. We do not, however, believe it would have




Figure 3 Number of studies analysing abiotic and biotic habitat variables by total spatial scale of study.

broadened the scope of the analyses presented as only five non-peer-reviewed articles appeared within our search criteria, three of which are known to have been published subsequently as peer-reviewed articles which were then included in our analysis. The exclusion of non-English peer-reviewed articles may well explain the low numbers of studies coming from South America and the complete lack of work originating from Africa and Asia. Nonetheless, we have no reason to believe that the patterns identified are likely to be geographically biased other than through the temperate restriction placed on the search criteria. The exclusion of articles prior to 1980 is not likely to have any significant bearing on our findings as we clearly demonstrate that work in this field did not develop substantially until post-1980, with only four articles published between 1980 and 1990. The tendency for authors to more readily report significant than non-significant relationships is likely to have increased the proportion of studies reporting significance. This publication bias is, however, expected to operate equally across all studies and spatial scales (Brett 1997) and therefore have little effect on our findings.

Spatial scale

The significance of scale in ecological investigations is well understood (Wiens 1989), and many studies have demonstrated the variable habitat use of temperate demersal fishes over a range of spatial and temporal scales (Bax *et al.* 1999; Lindholm *et al.* 1999; Walsh *et al.* 2004). Our analysis demonstrates that studies focusing on larger spatial scales primarily analyse abiotic habitat variables, whilst studies addressing biotic variables typically focus at smaller spatial scales. The tendency to study abiotic habitat variables over larger spatial scales may, however, not be wholly justified as our analysis showed no concomitant increase in the proportion of studies reporting significance at these larger spatial scales. As a result, our analyses indicate that there is no reason to limit the study of abiotic variables to large spatial scales. There was, however, no mismatch between the number of studies and the proportion of those studies in which biotic variables had a significant effect; both the total number of studies and the proportion of significant studies showed no pattern with spatial scale, except for the process of competition where




Figure 4 Number of non-mixed species studies (those analysing ≤ 4 species) reporting test statistics for each species and each abiotic habitat variable across the total spatial scales of study. Bars indicate the number of studies reporting significant (black) and non-significant (white) results. 'Other abiotic' includes the habitat variables of dissolved oxygen/level of hypoxia and light. (a) Depth, (b) Geography, (c) Hydrography, (d) Other abiotic, (e) Salinity, (f) Substrate, (g) Temperature, (h) Total abiotic (all abiotic habitat variables combined).

the total number of studies was very low. Thus, although the number of studies of biotic variables is in general low compared with the number of abiotic studies, the spread across all spatial scales for biotic habitat variables is likely justified.

The preponderance of large-scale abiotic studies could be a result of data availability, which will be discussed later. However, we believe that this also highlights the common assumption in this field that abiotic habitat variables are inherently linked to larger spatial scales and are most likely to influence the distribution of demersal fishes over larger spatial scales than do biotic habitat variables. This is, however, unlikely to be wholly the case, and numerous studies demonstrate

the importance of abiotic habitat variables over small spatial scales ($\leq 100 \text{ km}^2$) (Stoner *et al.* 2001; Laurel *et al.* 2009; Moore *et al.* 2009) and biotic habitat variables over larger spatial scales ($\geq 100 \text{ km}^2$) (Bjornstad *et al.* 1999; Vinagre *et al.* 2006; Le Pape *et al.* 2007). The lack of a relationship between the total spatial scale of study and the proportion of significant findings for the habitat variables analysed leads to the conclusion that there is no definitive spatial scale at which each of the different habitat variables significantly affect the distribution of temperate adult demersal fishes. It is therefore appropriate to analyse the effects of each habitat variable across all of the spatial scales analysed.




Figure 5 Number of non-mixed species studies (those analysing ≤ 4 species) reporting test statistics for each species and each biotic habitat variable across the total spatial scales of study. Bars indicate the number of studies reporting significant (black) and non-significant (white) results. (a) Biogenic complexity, (b) Competition, (c) Predation, (d) Prey resource, (e) Total biotic (all biotic habitat variables combined).




Figure 6 Number of studies carried out over different temporal scales: day = 0–24 h, month = 30 days, season = 90 days, annual = 365 days, na represents the studies that did not analyse any temporal pattern in fish abundance and/or distribution.

The high number of studies focussing on large spatial scales may also be explained by the origin of abiotic data. Almost half of the data used in studies

have originated from government-funded stock assessments and long-term, large spatial scale, scientific fish surveys. These data sources provide regular and large spatial scale information on fish abundance and local environmental variables that are directly comparable to the spatial scales of commercial fisheries. It is, however, important to consider that the large distances between sampling stations inherent with fishery-scale data sets mean that researchers are often unable to address the processes determining fish distribution that operate at smaller spatial scales (Hinz *et al.* 2006; Sullivan *et al.* 2006). An alternative but more costly approach, the use of multiscale, hierarchical surveys is likely to prove successful in elucidating the role of spatial scale in fish distribution (Pittman *et al.* 2004; Anderson and Yoklavich 2007; Anderson *et al.* 2009; Monk *et al.* 2011). These approaches warrant further investigation and development as there is still a paucity of studies using hierarchical designs in the field of demersal fish habitat determination.

Many studies utilizing large spatial scale data sets often take a 'data-mining' approach, searching for significant relationships between demersal fish abundances and measured habitat variables. This approach has been criticized for shifting our focus away from hypothesis-driven science, and potentially ascribing coincidental correlations or indirect relationships as direct causal links (Guisan *et al.* 2002). Insufficient knowledge of the multiple factors influencing the population dynamics of marine species makes it difficult to form prior assumptions about causal relationships. Consequently, correlative approaches that make few or even no prior assumptions about underlying causal relationships are considered a legitimate approach when attempting to understand the complex interactions between fish populations and their environment (see Valavanis *et al.* 2008). Correlations revealed from these approaches can in turn be used as the basis for subsequent hypothesis-driven studies aiming to determine demersal fish habitat requirements. New statistical approaches such as Quantile Regression (QR) and Structural Equation Modelling (SEM) allow better integration of sound ecological and statistical theory as well as the possibility of testing whether data are consistent with hypothesized causal pathways. Development of these approaches provides promising areas for future development (see Pittman *et al.* 2004; Vaz *et al.* 2008; Hermoso *et al.* 2011; Johnson *et al.* 2012).

Abiotic habitat variables

The bias towards focusing on abiotic variables may be related to the ease with which abiotic data are collected. With the exception of grain size analysis to determine sediment composition, all of the abiotic variables examined require relatively low sampling effort in comparison with that needed to investigate biotic variables (see Levin 1992). Abiotic habitat variables therefore provide a quick and often resource-effective means of collecting information on some of the habitat requirements that may determine the distribution and abundance of demersal fish species. Whether these variables and the relationships demonstrated are relevant to management strategies will likely be case dependent, determined by the species of interest, the habitat types covered and the spatial scales over which they occur.

Of the habitat variables studied, depth and substrate have received the most attention. Both may be

considered as useful proxies of other variables affecting the habitat choice of demersal fishes. Many previous studies note significant changes in community composition with depth and sediment type in benthic marine assemblages (Moranta *et al.* 1998; Kostylev *et al.* 2001; Hagberg *et al.* 2003). These relationships may indicate more complex associations with available prey communities and/or changes in habitat structure and complexity (Williams and Bax 2001; Anderson and Yoklavich 2007; Anderson *et al.* 2009). Although studies have successfully used depth, substrate and additional abiotic habitat variables to fill gaps in habitat knowledge (Last *et al.* 2010) and as surrogates for habitat suitability (Blanchard *et al.* 2005), this approach may be misleading depending on the scale of observation and the system under analysis (Chapman *et al.* 2010). Stevens and Connolly (2004) discuss the inadequacies of using surrogate abiotic measures to describe biotic communities at a range of spatial scales, concluding that less than 30% of the biological similarity between areas at scales covering tens of square kilometres could be explained using their abiotic proxies. Williams *et al.* (2009) similarly demonstrated that using a habitat characteristic describing geomorphic features over thousands of square kilometres to describe megafaunal communities led to a misrepresentation of species' rarities. It is therefore necessary that to describe and classify benthic communities and habitats with accuracy using only abiotic variables, they must first be calibrated against correlated biotic variables. This has been reflected in some habitat suitability studies that find that the best predictors of habitat suitability are models combining abiotic variables with biotic ones, such as prey density (Vinagre *et al.* 2006) and individual organismal traits (Le Pape *et al.* 2007). Planque *et al.* (2011) also discuss the inadequacy of expecting environmental (abiotic) habitat variables to fully explain the spatial distributions of fish populations unless their forcing effects are so strong as to override all other factors driving the distribution.

Biotic habitat variables

There is a general paucity of studies addressing the importance of biotic habitat variables in determining the distribution of demersal fish populations. Authors investigating the importance of different abiotic habitat variables often explained their results with reference to biotic habitat variables. Few studies, however, formally analysed the influence

of biotic habitat variables, particularly predation and competition, despite their known importance in determining the abundance and distribution of marine fishes (see Ward *et al.* 2006; Engelhard *et al.* 2008; Laurel *et al.* 2009). Studies addressing the importance of biotic habitat variables tended to focus at smaller total spatial scales of study compared with abiotic variables. Apart from substrate choice experiments, the two smallest spatial scale classes analysed (cm^2 and m^2) were dominated by biotic habitat variables. These spatial scales represent studies simulating environmental conditions using artificial laboratory habitat constructions. This approach provides the advantage of direct and continuous observation whilst controlling for additional habitat variables. The disadvantages, however, are well documented and highlighted by many authors working at this spatial scale. In an experiment aiming to investigate the role of variability in seafloor habitat structure on the survivorship of post-settlement juvenile Atlantic cod, Lindholm *et al.* (1999) acknowledged that limited space in such experiments does not allow for realistic foraging of predators or escape responses of prey. Manderson *et al.* (2000) also note that long experimental durations and small arena sizes provide increased predator-prey encounter rates in the laboratory, which are not wholly realistic for scaling up to conditions in the field.

Field validation of results from laboratory studies can help substantiate laboratory findings and may also elucidate previously overlooked variables important in determining habitat choice. Such validation may be undertaken directly by designing comparable experiments in the field or alternatively using pre-existing data to test relationships found within the laboratory. Stoner and Abookire (2002) provide an excellent example of the combination of laboratory and field experiments in their study of sediment preferences in Pacific halibut (*Hippoglossus stenolepis*, Pleuronectidae). Laboratory experiments showed detailed sediment preferences associated with fish size and burial capability. Subsequent field results overestimated the preferred sediment grain size of fish but still supported the hypothesis that sediment suitability was based on settlement capability. The authors also go on to discuss dissimilarities between laboratory and field results, highlighting discrepancies that may exist when extrapolating from small-scale manipulative studies to larger spatial scales.

Temporal aspects of habitat

Although the dynamic nature of fish habitat is widely acknowledged (Rice 2005), less than half of the studies reviewed looked at temporal patterns in habitat usage. Many studies utilizing data from long-term, large-scale stock assessment surveys grouped annual data, meaning that between-year differences in abundance and distribution were not considered. Pooling data may give wider applicability to findings but it also reduces the accuracy and hence predictive power of the results for the specific scenario that has been analysed (Valavanis *et al.* 2008). Differences in habitat suitability within and between habitats as well as important additional information such as fish year-class strength, variations in abundance and shifts in habitat use may be missed when temporal data are pooled. For example, studies by Manderson *et al.* (2002, 2003) on the growth, habitat variation and dynamic settlement in Winter flounder demonstrated that the influence of temperature and salinity on growth varied considerably from late spring into summer. Such studies also highlight the importance of considering location when analysing temporal differences in habitat suitability. In the previous example, freshwater inputs associated with estuaries caused complex changes in temperature and salinity. This is likely to cause large variations in predator-prey dynamics and the overall suitability of habitat important to the juvenile life stage of the study species (Gibson 1994; Manderson *et al.* 2006; Bachelier *et al.* 2009).

If researchers are to provide sound advice to managers on temporal distributions and predictions of future abundances, spatio-temporal approaches must be taken, in which temporal and spatial scales are matched *a priori*. For example, it would be impractical for a study covering thousands of square kilometres to attempt to analyse changes in habitat use over daily time scales. Similarly, over very small spatial scales, it would be inefficient to look at seasonal patterns in habitat use in demersal fish species, some of which are known to migrate large distances during annual spawning events (Armstrong *et al.* 1992; Hunter *et al.* 2003a). Such obvious examples highlight the clear need to plan sampling campaigns which incorporate appropriately matched temporal and spatial scales.

Ontogenetic changes in habitat use

Consideration of temporal patterns in habitat suitability is also important with respect to the ontogenetic changes which demersal fish species undergo during their development. Bachelier *et al.* (2009) showed how habitat use patterns of sub-adult red drum (*Sciaenops ocellatus*, Scianidae) in estuarine environments were age-dependent as well as region-dependent at large spatial scales, whilst Laurel *et al.* (2007) demonstrated habitat selection in Pacific halibut and rock sole (*Lepidopsetta polyxystra*, Pleuronectidae) was mediated by the interaction between temperature, ontogeny and density under laboratory conditions. Our analysis showed that fewer than half of studies clearly defined the life stages of the fish species analysed. Although combining juvenile and adult life stages may simplify sampling strategies and increase statistical power, it prevents ontogenetic changes in sensitivity and habitat requirements from being exposed. Those studies that did define life stage were often those focussed on post-larval settlement and habitat choices of juvenile fishes. Such investigations typically operate at small spatial scales, ranging from square centimetres to tens of square kilometres, often in laboratory or estuarine and shallow coastal environments (Stoner and Titgen 2003; Manderson *et al.* 2006; Laurel *et al.* 2007, 2009). To facilitate comparisons among studies and reduce ambiguity, there is a clear need for the specific life stages examined to be clarified. If neglected, conclusions may apply only to the dominant life stages within original samples and direct comparisons among studies may prove difficult.

Non-commercial and rare species

As well as a clear focus on mixed life stages and mixed species, studies have generally concentrated on commercially targeted demersal fish species. Few studies, however, have attempted to define the habitat variables determining the distribution and abundance of rare species, even where these are commercially valuable. For example, skate and ray species (Batoidea), which are known to be threatened by overfishing and habitat degradation (Hilton-Taylor 2010), have been seldom studied, even though such species often require species-specific studies to detect declines in population numbers (Dulvy *et al.* 2000). Such deficiencies in data and studies of rare species distributions may

hamper advice for conservation management strategies aiming to protect these or similar species.

Habitat suitability modelling

The number of studies in the field of demersal fish habitat associations has increased significantly over the past 20 years. This increased research effort has led to many advances in the methods and techniques used to address demersal fish habitat associations. Habitat suitability modelling has provided one attractive approach, as theoretically, results for a particular species should be applicable across systems (Rubec *et al.* 1999). Traditionally, many of these models assumed smooth, continuous and linear or simple polynomial relationships between habitat variables and fish populations. It is, however, often apparent that such functions cannot wholly predict the main processes modulating fish occurrence (see Caddy 2007). There now exist a suite of modelling approaches that are able to deal with more complex and biologically realistic relationships; GAMs combined with GIS platforms are generally considered the most well-developed method for modelling fish habitat use (Stoner *et al.* 2001, 2007; Valavanis *et al.* 2008; Bachelier *et al.* 2009; Katsanevakis and Maravelias 2009).

Progress may also be made through combinations of different models that account for weaknesses of each of the constituent models used. For example, machine-learning techniques are accurate predictors of complex nonlinear relationships with the additional ability to learn using training algorithms applied to random data subsets. They are, however, not good predictors of habitat suitability for new or unsampled sites (Maravelias *et al.* 2003; Pittman *et al.* 2009; Knudby *et al.* 2010). Presence-only algorithms on the other hand are able to assess new site suitabilities defined in terms of their environmental similarity (see Monk *et al.* 2010). Such algorithms could therefore potentially be integrated with machine-learning techniques to improve shortfalls in assessments of new site suitabilities whilst accurately predicting complex nonlinear relationships. Planque *et al.* (2011) also recommend a combinatory, multimodel approach to increase the accuracy of predictions and our overall understanding of factors controlling the spatial distribution of fish populations. Although an area showing promise, complex modelling approaches still need to address the numerous issues related to assumption violation, particularly

with respect to spatial auto-correlation and data independence as well as model validation, evaluation and the integration of sound ecological theory (Knudby *et al.* 2010).

Technological advancements

Technological developments have also played an important role in allowing researchers to answer increasingly complex and logistically challenging questions. Direct, *in situ* observations using underwater camera apparatus can inform us about patterns of distribution on small scales (Holmes *et al.* 2008), certain fish–habitat associations (Busby *et al.* 2005; Anderson and Yoklavich 2007) as well as detailed behavioural information (Stoner *et al.* 2008), all of which are unattainable using trawl methodologies alone. Similarly, advances in tagging technologies and in particular acoustic telemetry tags can give us information on locations and times of residency of individual fish (Lindholm *et al.* 2007; Alos *et al.* 2011; Andrews *et al.* 2011) as well as additional data such as depth, temperature and information on swimming behaviours (Hunter *et al.* 2003a,b). Such precise observations can give in-depth, real-time insights into the movement and behaviour of demersal fish and differential habitat uses when analysed in conjunction with corresponding habitat maps or benthic images (Sims *et al.* 2001).

Only 5–10% of the world's seafloor is currently mapped at resolutions similar to terrestrial studies (see Wright and Heyman 2008). This lack of benthic habitat maps over large scales has somewhat limited our ability to study the importance of landscape configuration and composition as well as benthic community structure on demersal fish distributions (Wright and Heyman 2008; Moore *et al.* 2011). Acoustic sensing devices are now widely recommended as a method to sample large areas of the benthic marine environment (Williams and Bax 2001). Sidescan sonars and multibeam swath bathymetry systems may now be used to produce accurate maps of seafloor substrates and bottom topography, allowing characterization of benthic habitats across large areas with potentially increased accuracy and decreased sampling times compared with sediment surveys and fishers' interpretations (Kloser *et al.* 2001; Kostylev *et al.* 2001; Freitas *et al.* 2006; Van der Kooij *et al.* 2011). However, the accuracy of a ground-truthed acoustic map, when extrapolated to larger scales, has often

been debated (Bax *et al.* 1999; Diaz *et al.* 2004; Brown *et al.* 2011). Such an approach for large-scale habitat classification has potential but will often require costly ground-truthing during field surveys (Freeman and Rogers 2003; Roberts *et al.* 2005; Holmes *et al.* 2008). The data density mismatches between physical and biological methods will, however, remain unsolved unless acoustic methods can routinely resolve the elusive biological components that make a physical substrate a habitat (Diaz *et al.* 2004). Until then, acoustic methods may still be used to better target benthic sampling, aid decisions regarding study scale or resolution, and provide good baselines from which more detailed habitat information can be determined (Freeman and Rogers 2003; Roberts *et al.* 2005).

A lot of work has been undertaken to relate acoustic maps with benthic communities, using a variety of ground-truthing methods in combination with multivariate analysis techniques (Walsh *et al.* 2004; Anderson and Yoklavich 2007; Holmes *et al.* 2008). Moore *et al.* (2010, 2011) provide a good example of the 'landscape' approach, relating benthic habitat to demersal fish distribution using a distance-based multivariate linear model (DISTLM). The authors found that a combination of depth and six of the 23 abiotic landscape indices explained 34.8% of the variation in the fish assemblage, demonstrating the validity of using broad-scale landscape analysis along with indices of landscape configuration and composition to explain distribution patterns in temperate demersal fish assemblages. Such approaches are now considered a vital first step in unravelling ecological complexities providing improved spatial information for management of marine systems (Brown *et al.* 2011).

Conclusion

Determining the habitat requirements of demersal fish species is inherently difficult because of the complex nature of marine ecosystems, the multiple factors affecting fish–habitat associations, the range of scales over which they act and the general difficulties of sampling marine habitats. Defining fish–habitat relationships will, however, be one of the necessary steps towards the advocated ecosystem-based approach to fisheries management and the sustainable exploitation of demersal fish stocks. Many developments in techniques and technologies show promise in elucidating the complex interac-

tions between demersal fishes and their habitat. The successful application of such developments will, however, strongly rely on the quality of the data used combined with understanding of the fundamental ecology of the systems and species under study (Austin 2007). Where possible, it will therefore be advantageous to test habitat variables under controlled experimental conditions (i.e. small-medium spatial scales), building results from such studies into larger scale models.

Our analysis highlights some important trends in the field of demersal fish habitat determination. The reasons behind the focus on larger scales are no doubt a result of a combination of factors. It is, however, clear that to advance the field, there should be a move towards the investigation of abiotic variables at smaller spatial scales as well as increased attention to the analysis of biotic habitat variables over all spatial scales of study. This will help describe distributions determined by abiotic habitat variables that may act over small spatial scales not previously considered and allow biotic-based causal relationships to be better explained. It will therefore be necessary to invest in the implementation of more, smaller spatial scale data collections or alternatively increase the resolution of larger spatial scale data sets. Work investigating the power of monitoring surveys to detect trends in abundance (e.g. Blanchard *et al.* 2008) will therefore prove invaluable in the design of future studies and surveys.

Defining temporal aspects of habitat will also prove valuable in advancing ecological understanding of the species under study. The inclusion of longer time scales and the consideration of temporal differences in habitat use may also provide important information on the cumulative effects of human-induced impacts, the overall status and recovery of impacted systems and increase capabilities to predict future change of the species or system under study (Hewitt *et al.* 2001). We argue that through attention to the areas highlighted herein, along with more holistic definitions of habitat, researchers are likely to be better equipped to inform management at a range of spatial and temporal scales.

Acknowledgements

The work presented in this paper benefited from discussions with E.H. Morgan, G.I. Lambert and J.O. Johnson. We are grateful to Paul Hart and two anonymous referees for their invaluable comments

on previous versions of this manuscript. A.F. Johnson was funded by the Natural Environment Research Council (UK).

References

- Abookire, A.A., Piatt, J.F. and Robards, M.D. (2000) Nearshore fish distributions in an Alaskan estuary in relation to stratification, temperature and salinity. *Estuarine Coastal and Shelf Science* **51**, 45–59.
- Alongi, D.M. (1998) Coastal ecosystem processes (Marine Science Series), CRC Press, Florida, US.
- Alos, J., March, D., Palmer, M., Grau, A. and Morales-Nin, B. (2011) Spatial and temporal patterns in *Serranus cabrilla* habitat use in the NW Mediterranean revealed by acoustic telemetry. *Marine Ecology-Progress Series* **427**, 173–186.
- Amara, R., Mahe, K., LePape, O. and Desroy, N. (2004) Growth, feeding and distribution of the solenette *Buglossidium luteum* with particular reference to its habitat preference. *Journal of Sea Research* **51**, 211–217.
- Amezcuca, F. and Nash, R.D.M. (2001) Distribution of the order Pleuronectiformes in relation to the sediment type in the North Irish Sea. *Journal of Sea Research* **45**, 293–301.
- Anderson, T.W. (2001) Predator responses, prey refuges, and density-dependent mortality of a marine fish. *Ecology* **82**, 245–257.
- Anderson, T.J. and Yoklavich, M.M. (2007) Multiscale habitat associations of deepwater demersal fishes off central California. *Fishery Bulletin* **105**, 168–179.
- Anderson, T.J., Syms, C., Roberts, D.A. and Howard, D.F. (2009) Multi-scale fish-habitat associations and the use of habitat surrogates to predict the organisation and abundance of deep-water fish assemblages. *Journal of Experimental Marine Biology and Ecology* **379**, 34–42.
- Andrews, K.S., Tolimieri, N., Williams, G.D., Samhour, J.F., Harvey, C.J. and Levin, P.S. (2011) Comparison of fine-scale acoustic monitoring systems using home range size of a demersal fish. *Marine Biology* **158**, 2377–2387.
- Araujo, F.G., de Azevedo, M.C.C., Silva, M.D., Pessanha, A.L.M., Gomes, I.D. and da Cruz, A.G. (2002) Environmental influences on the demersal fish assemblages in the Sepetiba Bay, Brazil. *Estuaries* **25**, 441–450.
- Ardron, J. (2008) The challenge of assessing whether the OSPAR network of marine protected areas is ecologically coherent. *Hydrobiologia* **606**, 45–53.
- Armstrong, J.D., Bagley, P.M. and Friede, I.G. (1992) Photographic and acoustic tracking observations of the behavior of the grenadier *Coryphaenoides (Nematonurus) armatus*, the eel *Synaphobranchus bathybius*, and other abyssal demersal fish in the North Atlantic Ocean. *Marine Biology* **112**, 535–544.
- Auster, P.J., Joy, K. and Valentine, P.C. (2001) Fish species and community distributions as proxies for seafloor

- habitat distributions: the Stellwagen Bank National Marine Sanctuary example (Northwest Atlantic, Gulf of Maine). *Environmental Biology of Fishes* **60**, 331–346.
- Auster, P.J., Lindholm, J., Schaub, S., Funnell, G., Kaufman, L.S. and Valentine, P.C. (2003) Use of sand wave habitats by silver hake. *Journal of Fish Biology* **62**, 143–152.
- Austin, M. (2007) Species distribution models and ecological theory: a critical assessment and some possible new approaches. *Ecological Modelling* **200**, 1–19.
- Bacheler, N.M., Paramore, L.M., Buckel, J.A. and Hightower, J.E. (2009) Abiotic and biotic factors influence the habitat use of an estuarine fish. *Marine Ecology-Progress Series* **377**, 263–277.
- Bax, N., Kloser, R., Williams, A., Gowlett-Holmes, K. and Ryan, T. (1999) Seafloor habitat definition for spatial management in fisheries: a case study on the continental shelf of southeast Australia. *Oceanologica Acta* **22**, 705–720.
- Belew, R. (2005) Scientific impact quantity and quality: analysis of two sources of bibliographic data. Vol. 2011. Arxiv preprint cs/0504036.
- Benaka, L. (1999) *Fish Habitat: Essential Fish Habitat and Rehabilitation*, Vol. 22, American Fisheries Society, Bethesda, MD.
- Bergstad, O.A., Menezes, G. and Hoines, A.S. (2008) Demersal fish on a mid-ocean ridge: distribution patterns and structuring factors. *Deep-Sea Research Part II-Topical Studies in Oceanography* **55**, 185–202.
- Biagi, F., Sartor, P., Ardizzone, G.D., Belcari, P., Belluscio, A. and Serena, F. (2002) Analysis of demersal assemblages of the Tuscany and Latium coasts (north-western Mediterranean). *Scientia Marina* **66**, 233–242.
- Bjornstad, O.N., Fromentin, J.M., Stenseth, N.C. and Gjosaeter, J. (1999) A new test for density-dependent survival: the case of coastal cod populations. *Ecology* **80**, 1278–1288.
- Blanchard, J.L., Mills, C., Jennings, S. *et al.* (2005) Distribution-abundance relationships for North Sea Atlantic cod (*Gadus morhua*): observation versus theory. *Canadian Journal of Fisheries and Aquatic Sciences* **62**, 2001–2009.
- Blanchard, J.L., Maxwell, D.L. and Jennings, S. (2008) Power of monitoring surveys to detect abundance trends in depleted populations: the effects of density-dependent habitat use, patchiness, and climate change. *ICES Journal of Marine Science* **65**, 111–120.
- Borg, A., Pihl, L. and Wennhage, H. (1997) Habitat choice by juvenile cod (*Gadus morhua* L.) on sandy soft bottoms with different vegetation types. *Helgolander Meeresuntersuchungen* **51**, 197–212.
- Brett, M.T. (1997) Meta-analysis in ecology. *Bulletin of the ecological society of America* **78**, 92–94.
- Brodeur, R.D. (2001) Habitat-specific distribution of Pacific ocean perch (*Sebastes alutus*) in Pribilof Canyon, Bering Sea. *Continental Shelf Research* **21**, 207–224.
- Brown, C.J., Smith, S.J., Lawton, P. and Anderson, J.T. (2011) Benthic habitat mapping: a review of progress towards improved understanding of the spatial ecology of the seafloor using acoustic techniques. *Estuarine Coastal and Shelf Science* **92**, 502–520.
- Busby, M.S., Mier, K.L. and Brodeur, R.D. (2005) Habitat associations of demersal fishes and crabs in the Pribilof Islands region of the Bering Sea. *Fisheries Research* **75**, 15–28.
- Caddy, J.F. (2007) *Marine Habitat and Cover: Their Importance for Productive Coastal Fishery Resources*, (Monographs on Oceanographic Methodology, Vol. 11, UNESCO Publishing, Paris.
- Cartes, J.E., Hidalgo, M., Papiol, V., Massuti, E. and Moranta, J. (2009) Changes in the diet and feeding of the hake *Merluccius merluccius* at the shelf-break of the Balearic Islands: influence of the mesopelagic-boundary community. *Deep-Sea Research Part I-Oceanographic Research Papers* **56**, 344–365.
- Chapman, M., Tolhurst, T., Murphy, R. and Underwood, A. (2010) Complex and inconsistent patterns of variation in benthos, micro-algae and sediment over multiple spatial scales. *Marine Ecology Progress Series* **398**, 33–47.
- Chatfield, B.S., Van Niel, K.P., Kendrick, G.A. and Harvey, E.S. (2010) Combining environmental gradients to explain and predict the structure of demersal fish distributions. *Journal of Biogeography* **37**, 593–605.
- Cincotta, R., Wisniewski, J. and Engelman, R. (2000) Human population in the biodiversity hotspots. *Nature* **404**, 990–992.
- Colloca, F., Carpentieri, P., Balestri, E. and Ardizzone, G. (2010) Food resource partitioning in a Mediterranean demersal fish assemblage: the effect of body size and niche width. *Marine Biology* **157**, 565–574.
- Damalas, D., Maravelias, C.D., Katsanevakis, S., Karageorgis, A.R. and Papaconstantinou, C. (2010) Seasonal abundance of non-commercial demersal fish in the eastern Mediterranean Sea in relation to hydrographic and sediment characteristics. *Estuarine Coastal and Shelf Science* **89**, 107–118.
- De Ben, W.A., Clothier, W.D., Ditsworth, G.R. and Baumgartner, D.J. (1990) Spatio-temporal fluctuations in the distribution and abundance of demersal fish and epibenthic crustaceans in Yaquina Bay, Oregon. *Estuaries* **13**, 469–478.
- Dean, T.A., Haldorson, L., Laur, D.R., Jewett, S.C. and Blanchard, A. (2000) The distribution of nearshore fishes in kelp and eelgrass communities in Prince William Sound, Alaska: associations with vegetation and physical habitat characteristics. *Environmental Biology of Fishes* **57**, 271–287.
- Demestre, M., Sanchez, P. and Abello, P. (2000) Demersal fish assemblages and habitat characteristics on the continental shelf and upper slope of the north-western Mediterranean. *Journal of the Marine Biological Association of the United Kingdom* **80**, 981–988.
- Diaz, R.J., Solan, M. and Valente, R.M. (2004) A review of approaches for classifying benthic habitats and evaluat-

- ing habitat quality. *Journal of Environmental Management* **73**, 165–181.
- Dulvy, N.K., Metcalfe, J.D., Glanville, J., Pawson, M.G. and Reynolds, J.D. (2000) Fishery stability, local extinctions, and shifts in community structure in skates. *Conservation Biology* **14**, 283–293.
- Eastwood, P.D., Meaden, G.J., Carpentier, A. and Rogers, S.I. (2003) Estimating limits to the spatial extent and suitability of sole (*Solea solea*) nursery grounds in the Dover Strait. *Journal of Sea Research* **50**, 151–165.
- Eby, L.A., Crowder, L.B., McClellan, C.M., Peterson, C.H. and Powers, M.J. (2005) Habitat degradation from intermittent hypoxia: impacts on demersal fishes. *Marine Ecology-Progress Series* **291**, 249–261.
- Edgar, G.J. and Shaw, C. (1995) The production and trophic ecology of shallow-water fish assemblages in southern Australia (3) General relationships between sediments, seagrasses, invertebrates and fishes. *Journal of Experimental Marine Biology and Ecology* **194**, 107–131.
- Ehrich, S., Stelzenmuller, V. and Adlerstein, S. (2009) Linking spatial pattern of bottom fish assemblages with water masses in the North Sea. *Fisheries Oceanography* **18**, 36–50.
- Ekau, W. (1990) Demersal fish fauna of the Weddell Sea, Antarctica. *Antarctic Science* **2**, 129–137.
- Ellis, J.R., Rogers, S.I. and Freeman, S.M. (2000) Demersal assemblages in the Irish Sea, St George's Channel and Bristol Channel. *Estuarine Coastal and Shelf Science* **51**, 299–315.
- Engelhard, G.H., Van der Kooij, J., Bell, E.D. et al. (2008) Fishing mortality versus natural predation on diurnally migrating sandeels *Ammodytes marinus*. *Marine Ecology-Progress Series* **369**, 213–227.
- Farina, A.C., Freire, J. and GonzalezGurriaran, E. (1997) Demersal fish assemblages in the Galician continental shelf and upper slope (NW Spain): spatial structure and long-term changes. *Estuarine Coastal and Shelf Science* **44**, 435–454.
- Florentino, F., Badalamenti, F., D'Anna, G. et al. (2008) Changes in spawning-stock structure and recruitment pattern of red mullet, *Mullus barbatus*, after a trawl ban in the Gulf of Castellammare (central Mediterranean Sea). *ICES Journal of Marine Science* **65**, 1175–1183.
- Folke, C., Carpenter, S., Walker, B. et al. (2004) Regime shifts, resilience, and biodiversity in ecosystem management. *Annual Review of Ecology Evolution and Systematics* **35**, 557–581.
- Francis, M.P., Hurst, R.J., McArdle, B.H., Bagley, N.W. and Anderson, O.F. (2002) New Zealand demersal fish assemblages. *Environmental Biology of Fishes* **65**, 215–234.
- Frank, K.T., Petrie, B., Choi, J.S. and Leggett, W.C. (2005) Trophic cascades in a formerly cod-dominated ecosystem. *Science* **308**, 1621–1623.
- Freeman, S.M. and Rogers, S.I. (2003) A new analytical approach to the characterisation of macro-epibenthic habitats: linking species to the environment. *Estuarine Coastal and Shelf Science* **56**, 749–764.
- Freitas, R., Sampalo, L., Oliveira, J., Rodrigues, A.M. and Quintino, V. (2006) Validation of soft bottom benthic habitats identified by single-beam acoustics. *Marine Pollution Bulletin* **53**, 72–79.
- Frid, C., Paramor, O. and Scott, C. (2005) Ecosystem-based fisheries management: progress in the NE Atlantic. *Marine Policy* **29**, 461–469.
- Fromentin, J.M., Myers, R.A., Bjornstad, O.N., Stenseth, N.C., Gjosaeter, J. and Christie, H. (2001) Effects of density-dependent and stochastic processes on the regulation of cod populations. *Ecology* **82**, 567–579.
- Frost, H. and Andersen, P. (2006) The common fisheries policy of the European union and fisheries economics. *Marine Policy* **30**, 737–746.
- Fujita, T., Inada, T. and Ishito, Y. (1995) Depth-gradient structure of the demersal fish community on the continental-shelf and upper slope off Sendai Bay, Japan. *Marine Ecology-Progress Series* **118**, 13–23.
- Gabriel, W.L. (1992) Persistence of demersal fish assemblages between Cape Hatteras and Nova Scotia, north-west Atlantic. *Journal of Northwest Atlantic Fishery Science* **14**, 29–46.
- Gaertner, J.C., Mazouni, N., Sabatier, R. and Millet, B. (1999) Spatial structure and habitat associations of demersal assemblages in the Gulf of Lions: a multi-compartmental approach. *Marine Biology* **135**, 199–208.
- Gaines, S.D., White, C., Carr, M.H. and Palumbi, S.R. (2010) Designing marine reserve networks for both conservation and fisheries management. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 18286–18293.
- Gelcich, S., Defeo, O., Iribarne, O. et al. (2009) Marine ecosystem-based management in the Southern Cone of South America: stakeholder perceptions and lessons for implementation. *Marine Policy* **33**, 801–806.
- Gibson, R.N. (1994) Impact of habitat quality and quantity on the recruitment of juvenile flatfishes. *Netherlands Journal of Sea Research* **32**, 191–206.
- Gibson, R. and Ezzi, I. (1987) Feeding relationships of a demersal fish assemblage on the west coast of Scotland. *Journal of Fish Biology* **31**, 55–69.
- Gibson, R.N. and Robb, L. (1992) The relationship between body size, sediment grain-size and the burying ability of juvenile plaice, *Pleuronectes platessa* L. *Journal of Fish Biology* **40**, 771–778.
- Guisan, A., Edwards, T.C. and Hastie, T. (2002) Generalized linear and generalized additive models in studies of species distributions: setting the scene. *Ecological Modelling* **157**, 89–100.
- Gutt, J. and Ekau, W. (1996) Habitat partitioning of dominant high Antarctic demersal fish in the Weddell Sea and Lazarev Sea. *Journal of Experimental Marine Biology and Ecology* **206**, 25–37.

- Hagberg, J., Jonzen, N., Lundberg, P. and Ripa, J. (2003) Uncertain biotic and abiotic interactions in benthic communities. *Oikos* **100**, 353–361.
- Harzing, A.W. and Van der Wal, R. (2008) Google Scholar as a new source for citation analysis. *Ethics in Science and Environmental politics* **8**, 61–73.
- Heenan, A., Simpson, S.D., Meekan, M.G., Healy, S.D. and Braithwaite, V.A. (2009) Restoring depleted coral-reef fish populations through recruitment enhancement: a proof of concept. *Journal of Fish Biology* **75**, 1857–1867.
- Hermoso, V., Clavero, M. and Kennard, M.J. (2011) Determinants of fine-scale homogenization and differentiation of native freshwater fish faunas in a Mediterranean Basin: implications for conservation. *Diversity and Distributions* **18**, 1–12.
- Hewitt, J.E., Thrush, S.E. and Cummings, V.J. (2001) Assessing environmental impacts: effects of spatial and temporal variability at likely impact scales. *Ecological Applications* **11**, 1502–1516.
- Hiddink, J., Jennings, S., Kaiser, M., Queirós, A., Duplisea, D. and Piet, G. (2006) Cumulative impacts of seabed trawl disturbance on benthic biomass, production, and species richness in different habitats. *Canadian Journal of Fisheries and Aquatic Sciences* **63**, 721–736.
- Hiddink, J.G., Johnson, A.F., Kingham, R. and Hinz, H. (2011) Could our fisheries be more productive? Indirect negative effects of bottom trawl fisheries on fish condition. *Journal of Applied Ecology* **48**, 1441–1449.
- Hilton-Taylor, C. (2010) *IUCN Red List of Threatened Species. Version 2010.1, 2011.1*, <http://www.iucnredlist.org>. Downloaded on 24 July 2011
- Hindell, J.S., Jenkins, G.P. and Keough, M.J. (2001) Spatial and temporal variability in the effects of fish predation on macrofauna in relation to habitat complexity and cage effects. *Marine Ecology-Progress Series* **224**, 231–250.
- Hinz, H., Kaiser, M.J., Bergmann, M., Rogers, S.I. and Armstrong, M.J. (2003) Ecological relevance of temporal stability in regional fish catches. *Journal of Fish Biology* **63**, 1219–1234.
- Hinz, H., Kroncke, I. and Ehrich, S. (2005) The feeding strategy of dab *Limanda limanda* in the southern North Sea: linking stomach contents to prey availability in the environment. *Journal of Fish Biology* **67**, 125–145.
- Hinz, H., Bergmann, M., Shucksmith, R., Kaiser, M.J. and Rogers, S.I. (2006) Habitat association of plaice, sole, and lemon sole in the English Channel. *ICES Journal of Marine Science* **63**, 912–927.
- Hinz, H., Prieto, V. and Kaiser, M.J. (2009) Trawl disturbance on benthic communities: chronic effects and experimental predictions. *Ecological Applications* **19**, 761–773.
- Holmes, K.W., Van Niel, K.P., Radford, B., Kendrick, G.A. and Grove, S.L. (2008) Modelling distribution of marine benthos from hydroacoustics and underwater video. *Continental Shelf Research* **28**, 1800–1810.
- Hughes, J.E., Deegan, L.A., Wyda, J.C., Weaver, M.J. and Wright, A. (2002) The effects of eelgrass habitat loss on estuarine fish communities of southern new England. *Estuaries* **25**, 235–249.
- Hunter, E., Aldridge, J.N., Metcalfe, J.D. and Arnold, G.P. (2003a) Geolocation of free-ranging fish on the European continental shelf as determined from environmental variables – I. Tidal location method. *Marine Biology* **142**, 601–609.
- Hunter, E., Metcalfe, J.D. and Reynolds, J.D. (2003b) Migration route and spawning area fidelity by North Sea plaice. *Proceedings of the Royal Society of London Series B-Biological Sciences* **270**, 2097–2103.
- Hyndes, G., Platell, M., Potter, I. and Lenanton, R. (1999) Does the composition of the demersal fish assemblages in temperate coastal waters change with depth and undergo consistent seasonal changes? *Marine Biology* **134**, 335–352.
- Iglesias, J. (1981) Spatial and temporal changes in the demersal fish community of the Ria de Arosa (NW Spain). *Marine Biology* **65**, 199–208.
- Jackson, J.B.C., Kirby, M.X., Berger, W.H. *et al.* (2001) Historical overfishing and the recent collapse of coastal ecosystems. *Science* **293**, 629–638.
- Johnson, S.W., Murphy, M.L. and Csepp, D.J. (2003) Distribution, habitat, and behavior of rockfishes, *Sebastes* spp., in nearshore waters of southeastern Alaska: observations from a remotely operated vehicle. *Environmental Biology of Fishes* **66**, 259–270.
- Johnson, A.F., Valls, M., Moranta, M., Jenkins, S.R., Hiddink, J.G. and Hinz, H. (2012) Effect of prey abundance and size on the distribution of demersal fishes. *Canadian Journal of Fisheries and Aquatic Sciences* **69**, 191–200.
- Kaiser, M. (1998) Significance of bottom fishing disturbance. *Conservation Biology* **12**, 1230–1235.
- Kaiser, M.J., Collie, J.S., Hall, S.J., Jennings, S. and Poiner, I.R. (2002) Modification of marine habitats by trawling activities: prognosis and solutions. *Fish and Fisheries* **3**, 114–136.
- Kaiser, M.J., Bergmann, M., Hinz, H. *et al.* (2004) Demersal fish and epifauna associated with sandbank habitats. *Estuarine Coastal and Shelf Science* **60**, 445–456.
- Karagiannakos, A. (1996) Total allowable catch (TAC) and quota management system in the European union. *Marine Policy* **20**, 235–248.
- Katsanevakis, S. and Maravelias, C.D. (2009) Bathymetric distribution of demersal fish in the Aegean and Ionian Seas based on generalized additive modeling. *Fisheries Science* **75**, 13–23.
- Katsanevakis, S., Maravelias, C.D., Damalas, D. *et al.* (2009) Spatiotemporal distribution and habitat use of commercial demersal species in the eastern Mediterranean Sea. *Fisheries Oceanography* **18**, 439–457.
- Keller, A.A., Simon, V., Chan, F. *et al.* (2010) Demersal fish and invertebrate biomass in relation to an offshore

- hypoxic zone along the US West Coast. *Fisheries Oceanography* **19**, 76–87.
- Kloser, R.J., Bax, N.J., Ryan, T., Williams, A. and Barker, B.A. (2001) Remote sensing of seabed types in the Australian South East Fishery; development and application of normal incident acoustic techniques and associated 'ground truthing'. *Marine and Freshwater Research* **52**, 475–489.
- Knudby, A., Brenning, A. and LeDrew, E. (2010) New approaches to modelling fish-habitat relationships. *Ecological Modelling* **221**, 503–511.
- Koslow, J.A., Bulman, C.M. and Lyle, J.M. (1994) The mid-slope demersal fish community off southeastern Australia. *Deep-Sea Research Part I-Oceanographic Research Papers* **41**, 113–141.
- Koss, R., Bellgrove, A., Ierodionou, D., Gilmour, P. and Bunce, A. (2005) Sea search: community-based monitoring of Victoria's Marine National Parks and Marine Sanctuaries: subtidal reef monitoring. *Parks Victoria Technical Series* **1**, 8.
- Kostylev, V.E., Todd, B.J., Fader, G.B.J., Courtney, R.C., Cameron, G.D.M. and Pickrill, R.A. (2001) Benthic habitat mapping on the Scotian shelf based on multi-beam bathymetry, surficial geology and sea floor photographs. *Marine Ecology-Progress Series* **219**, 121–137.
- Laidig, T.E., Watters, D.L. and Yoklavich, M.M. (2009) Demersal fish and habitat associations from visual surveys on the central California shelf. *Estuarine Coastal and Shelf Science* **83**, 629–637.
- Langhamer, O. and Wilhelmsson, D. (2009) Colonisation of fish and crabs of wave energy foundations and the effects of manufactured holes – a field experiment. *Marine Environmental Research* **68**, 151–157.
- Last, P.R., Lyne, V.D., Williams, A., Davies, C.R., Butler, A.J. and Yearsley, G.K. (2010) A hierarchical framework for classifying seabed biodiversity with application to planning and managing Australia's marine biological resources. *Biological Conservation* **143**, 1675–1686.
- Laurel, B.J., Stoner, A.W. and Hurst, T.P. (2007) Density-dependent habitat selection in marine flatfish: the dynamic role of ontogeny and temperature. *Marine Ecology-Progress Series* **338**, 183–192.
- Laurel, B.J., Ryer, C.H., Knoth, B. and Stoner, A.W. (2009) Temporal and ontogenetic shifts in habitat use of juvenile Pacific cod (*Gadus macrocephalus*). *Journal of Experimental Marine Biology and Ecology* **377**, 28–35.
- Lazzari, M.A. and Tupper, B. (2002) Importance of shallow water habitats for demersal fishes and decapod crustaceans in Penobscot Bay, Maine. *Environmental Biology of Fishes* **63**, 57–66.
- Le Pape, O., Chauvet, F., Mahevas, S., Lazure, P., Guerault, D. and Desaunay, Y. (2003) Quantitative description of habitat suitability for the juvenile common sole (*Solea solea*, L.) in the Bay of Biscay (France) and the contribution of different habitats to the adult population. *Journal of Sea Research* **50**, 139–149.
- Le Pape, O., Baulier, L., Cloarec, A., Martin, J., Le Loc'h, F. and Desaunay, Y. (2007) Habitat suitability for juvenile common sole (*Solea solea*, L.) in the Bay of Biscay (France): a quantitative description using indicators based on epibenthic fauna. *Journal of Sea Research* **57**, 126–136.
- Leathwick, J.R., Elith, J., Francis, M.P., Hastie, T. and Taylor, P. (2006) Variation in demersal fish species richness in the oceans surrounding New Zealand: an analysis using boosted regression trees. *Marine Ecology-Progress Series* **321**, 267–281.
- Levin, S.A. (1992) The problem of pattern and scale in ecology. *Ecology* **73**, 1943–1967.
- Levin, P.S. (1994) Small-scale recruitment variation in a temperate fish – the roles of macrophytes and food-supply. *Environmental Biology of Fishes* **40**, 271–281.
- Lindholm, J.B., Auster, P.J. and Kaufman, L.S. (1999) Habitat-mediated survivorship of juvenile (0-year) Atlantic cod *Gadus morhua*. *Marine Ecology-Progress Series* **180**, 247–255.
- Lindholm, J.B., Auster, P.J., Ruth, M. and Kaufman, L. (2001) Modeling the effects of fishing and implications for the design of marine protected areas: Juvenile fish responses to variations in seafloor habitat. *Conservation Biology* **15**, 424–437.
- Lindholm, J., Auster, P.J. and Knight, A. (2007) Site fidelity and movement of adult Atlantic cod *Gadus morhua* at deep boulder reefs in the western Gulf of Maine, USA. *Marine Ecology-Progress Series* **342**, 239–247.
- Link, J.S. (2002) What does ecosystem-based fisheries management mean? *Fisheries* **27**, 18–21.
- Logerwell, E.A., Aydin, K., Barbeaux, S. et al. (2005) Geographic patterns in the demersal ichthyofauna of the Aleutian Islands. *Fisheries Oceanography* **14**, 93–112.
- Macpherson, E. (1981) Resource partitioning in a Mediterranean demersal fish community. *Marine Ecology Progress Series* **4**, 183–193.
- Magnussen, E. (2002) Demersal fish assemblages of Faroe Bank: species composition, distribution, biomass spectrum and diversity. *Marine Ecology-Progress Series* **238**, 211–225.
- Mahon, R. and Smith, R.W. (1989) Demersal fish assemblages on the Scotian Shelf, northwest Atlantic – spatial-distribution and persistence. *Canadian Journal of Fisheries and Aquatic Sciences* **46**, 134–152.
- Manderson, J.P., Phelan, B.A., Stoner, A.W. and Hilbert, J. (2000) Predator-prey relations between age-1 + summer flounder (*Paralichthys dentatus*, Linnaeus) and age-0 winter flounder (*Pseudopleuronectes americanus*, Walbaum): predator diets, prey selection, and effects of sediments and macrophytes. *Journal of Experimental Marine Biology and Ecology* **251**, 17–39.
- Manderson, J.P., Phelan, B.A., Meise, C. et al. (2002) Spatial dynamics of habitat suitability for the growth of

- newly settled winter flounder *Pseudopleuronectes americanus* in an estuarine nursery. *Marine Ecology-Progress Series* **228**, 227–239.
- Manderson, J.P., Pessutti, J., Meise, C., Johnson, D. and Shaheen, P. (2003) Winter flounder settlement dynamics and the modification of settlement patterns by post-settlement processes in a NW Atlantic estuary. *Marine Ecology-Progress Series* **253**, 253–267.
- Manderson, J.P., Pessutti, J., Shaheen, P. and Juanes, F. (2006) Dynamics of early juvenile winter flounder predation risk on a North West Atlantic estuarine nursery ground. *Marine Ecology-Progress Series* **328**, 249–265.
- Maravelias, C.D., Haralabous, J. and Papaconstantinou, C. (2003) Predicting demersal fish species distributions in the Mediterranean Sea using artificial neural networks. *Marine Ecology-Progress Series* **255**, 249–258.
- Massuti, E. and Renones, O. (2005) Demersal resource assemblages in the trawl fishing grounds off the Balearic Islands (western Mediterranean). *Scientia Marina* **69**, 167–181.
- Maxwell, D.L., Stelzenmuller, V., Eastwood, P.D. and Rogers, S.I. (2009) Modelling the spatial distribution of plaice (*Pleuronectes platessa*), sole (*Solea solea*) and thornback ray (*Raja clavata*) in UK waters for marine management and planning. *Journal of Sea Research* **61**, 258–267.
- McClatchie, S., Millar, R.B., Webster, F., Lester, P.J., Hurst, R. and Bagley, N. (1997) Demersal fish community diversity off New Zealand: is it related to depth, latitude and regional surface phytoplankton? *Deep-Sea Research Part I-Oceanographic Research Papers* **44**, 647–667.
- McConnaughey, R.A. and Smith, K.R. (2000) Associations between flatfish abundance and surficial sediments in the eastern Bering Sea. *Canadian Journal of Fisheries and Aquatic Sciences* **57**, 2410–2419.
- McCormick, M.I. (1998) Ontogeny of diet shifts by a microcarnivorous fish, *Cheilodactylus spectabilis*: relationship between feeding mechanics, microhabitat selection and growth. *Marine Biology* **132**, 9–20.
- Medina, A., Brethes, J.C., Sevigny, J.M. and Zakardjian, B. (2007) How geographic distance and depth drive ecological variability and isolation of demersal fish communities in an archipelago system (Cape Verde, Eastern Atlantic Ocean). *Marine Ecology-an Evolutionary Perspective* **28**, 404–417.
- Meho, L.I. and Yang, K. (2007) Impact of data sources on citation counts and rankings of LIS faculty: web of science versus scopus and google scholar. *Journal of the American Society for Information Science and Technology* **58**, 2105–2125.
- Methven, D.A., Haedrich, R.L. and Rose, G.A. (2001) The fish assemblage of a Newfoundland estuary: diel, monthly and annual variation. *Estuarine Coastal and Shelf Science* **52**, 669–687.
- Monk, J., Ierodiaconou, D., Versace, V.L. et al. (2010) Habitat suitability for marine fishes using presence-only modelling and multibeam sonar. *Marine Ecology-Progress Series* **420**, 157–174.
- Monk, J., Ierodiaconou, D., Bellgrove, A., Harvey, E. and Laurenson, L. (2011) Remotely sensed hydroacoustics and observation data for predicting fish habitat suitability. *Continental Shelf Research* **31**, S17–S27.
- Moore, C.H., Harvey, E.S. and Van Niel, K.P. (2009) Spatial prediction of demersal fish distributions: enhancing our understanding of species-environment relationships. *Ices Journal of Marine Science* **66**, 2068–2075.
- Moore, C.H., Harvey, E.S. and Van Niel, K. (2010) The application of predicted habitat models to investigate the spatial ecology of demersal fish assemblages. *Marine Biology* **157**, 2717–2729.
- Moore, C.H., Van Niel, K. and Harvey, E.S. (2011) The effect of landscape composition and configuration on the spatial distribution of temperate demersal fish. *Ecography* **34**, 425–435.
- Moranta, J., Stefanescu, C., Massuti, E., Morales-Nin, B. and Lloris, D. (1998) Fish community structure and depth-related trends on the continental slope of the Balearic Islands (Algerian basin, western Mediterranean). *Marine Ecology-Progress Series* **171**, 247–259.
- Murawski, S.A., Brown, R., Lai, H.L., Rago, P.J. and Hendrickson, L. (2000) Large-scale closed areas as a fishery-management tool in temperate marine systems: the Georges bank experience. *Bulletin of Marine Science* **66**, 775–798.
- Nicolas, D., Le Loc'h, F., Desaunay, Y., Hamon, D., Blanchet, A. and Le Pape, O. (2007) Relationships between benthic macrofauna and habitat suitability for juvenile common sole (*Solea solea*, L.) in the Vilaine estuary (Bay of Biscay, France) nursery ground. *Estuarine Coastal and Shelf Science* **73**, 639–650.
- Ordines, F. and Massuti, E. (2009) Relationships between macro-epibenthic communities and fish on the shelf grounds of the western Mediterranean. *Aquatic Conservation-Marine and Freshwater Ecosystems* **19**, 370–383.
- Pauly, D. and Stergiou, K.I. (2005) Equivalence of results from two citation analyses: thomson's ISI citation index and Google's Scholar service. *Ethics in Science and Environmental Politics*, **3**, 3–35.
- Pauly, D., Watson, R. and Alder, J. (2005) Global trends in world fisheries: impacts on marine ecosystems and food security. *Philosophical Transactions of the Royal Society B-Biological Sciences* **360**, 5–12.
- Perry, A., Low, P., Ellis, J. and Reynolds, J. (2005) Climate change and distribution shifts in marine fishes. *Science* **308**, 1912.
- Persohn, C., Lorange, P. and Trenkel, V.M. (2009) Habitat preferences of selected demersal fish species in the Bay of Biscay and Celtic Sea, North-East Atlantic. *Fisheries Oceanography* **18**, 268–285.

- Phelan, B.A., Manderson, J.P., Stoner, A.W. and Bejda, A.J. (2001) Size-related shifts in the habitat associations of young-of-the-year winter flounder (*Pseudopleuronectes americanus*): field observations and laboratory experiments with sediments and prey. *Journal of Experimental Marine Biology and Ecology* **257**, 297–315.
- Piet, G.J., Pfisterer, A.B. and Rijnsdorp, A.D. (1998) On factors structuring the flatfish assemblage in the southern North Sea. *Journal of Sea Research* **40**, 143–152.
- Pihl, L., Baden, S.P. and Diaz, R.J. (1991) Effects of periodic hypoxia on distribution of demersal fish and crustaceans. *Marine Biology* **108**, 349–360.
- Pikitch, E.K., Santora, C., Babcock, E.A. et al. (2004) Ecosystem-based fishery management. *Science* **305**, 346–347.
- Pittman, S.J., McAlpine, C.A. and Pittman, K.M. (2004) Linking fish and prawns to their environment: a hierarchical landscape approach. *Marine Ecology-Progress Series* **283**, 233–254.
- Pittman, S.J., Costa, B.M. and Battista, T.A. (2009) Using lidar bathymetry and boosted regression trees to predict the diversity and abundance of fish and corals. *Journal of Coastal Research* **25**, 27–38.
- Planque, B., Loots, C., Petitgas, P., Lindstrom, U. and Vaz, S. (2011) Understanding what controls the spatial distribution of fish populations using a multi-model approach. *Fisheries Oceanography* **20**, 1–17.
- Queiros, A.M., Hiddink, J.G., Kaiser, M.J. and Hinz, H. (2006) Effects of chronic bottom trawling disturbance on benthic biomass, production and size spectra in different habitats. *Journal of Experimental Marine Biology and Ecology* **335**, 91–103.
- Quiroga, E., Sellanes, J., Arntz, W.E., Gerdes, D., Gallardo, V.A. and Hebbeln, D. (2009) Benthic megafaunal and demersal fish assemblages on the Chilean continental margin: the influence of the oxygen minimum zone on bathymetric distribution. *Deep-Sea Research Part II Topical Studies in Oceanography* **56**, 1112–1123.
- Reiss, H., Degraer, S., Duineveld, G.C.A. et al. (2010) Spatial patterns of infauna, epifauna, and demersal fish communities in the North Sea. *ICES Journal of Marine Science* **67**, 278–293.
- Rice, J.C. (2005) Understanding fish habitat ecology to achieve conservation. *Journal of Fish Biology* **67**, 1–22.
- Roberts, C.M. and Ormond, R.F.G. (1987) Habitat complexity and coral-reef fish diversity and abundance on Red-Sea fringing reefs. *Marine Ecology-Progress Series* **41**, 1–8.
- Roberts, J.M., Brown, C.J., Long, D. and Bates, C.R. (2005) Acoustic mapping using a multibeam echosounder reveals cold-water coral reefs and surrounding habitats. *Coral Reefs* **24**, 654–669.
- Robichaud, D. and Rose, G.A. (2006) Density-dependent distribution of demersal juvenile Atlantic cod (*Gadus morhua*) in Placentia Bay, Newfoundland. *ICES Journal of Marine Science* **63**, 766–774.
- Rubec, P.J., Bexley, J.C.W., Norris, H. et al. (1999) Suitability modeling to delineate habitat essential to sustainable fisheries. *Fish Habitat: Essential Fish Habitat and Rehabilitation* **22**, 108–133.
- Ryer, C.H., Stoner, A.W. and Titgen, R.H. (2004) Behavioral mechanisms underlying the refuge value of benthic habitat structure for two flatfishes with differing anti-predator strategies. *Marine Ecology-Progress Series* **268**, 231–243.
- Ryer, C.H., Stoner, A.W., Spencer, M.L. and Abookire, A.A. (2007) Presence of larger flatfish modifies habitat preference by Age-0 northern rock sole *Lepidopsetta polyxystra*. *Marine Ecology-Progress Series* **342**, 227–238.
- Seaman, W. (2008) *Coastal Artificial Habitats for Fishery and Environmental Management and Scientific Advancement*, (Fisheries for global welfare and environment. Memorial book of the 5th world Fisheries Congress 2008., Vol. 1, Terrapub, Tokyo.
- Sedberry, G.R. and Vandolah, R.F. (1984) Demersal fish assemblages associated with hard bottom habitat in the South-Atlantic Bight of the USA. *Environmental Biology of Fishes* **11**, 241–258.
- Selleslagh, J. and Amara, R. (2008) Environmental factors structuring fish composition and assemblages in a small macrotidal estuary (Eastern English Channel). *Estuarine Coastal and Shelf Science* **79**, 507–517.
- Shephard, S., Brophy, D. and Reid, D.G. (2010) Can bottom trawling indirectly diminish carrying capacity in a marine ecosystem? *Marine Biology (Berlin)* **157**, 2375–2381.
- Shepherd, T.D. and Litvak, M.K. (2004) Density-dependent habitat selection and the ideal free distribution in marine fish spatial dynamics: considerations and cautions. *Fish and Fisheries* **5**, 141–152.
- Shucksmith, R., Hinz, H., Bergmann, M. and Kaiser, M.J. (2006) Evaluation of habitat use by adult plaice (*Pleuronectes platessa* L.) using underwater video survey techniques. *Journal of Sea Research* **56**, 317–328.
- Simpson, M.R. and Walsh, S.J. (2004) Changes in the spatial structure of Grand Bank yellowtail flounder: testing MacCall's basin hypothesis. *Journal of Sea Research* **51**, 199–210.
- Sims, D.W., Nash, J.P. and Morritt, D. (2001) Movements and activity of male and female dogfish in a tidal sea lough: alternative behavioural strategies and apparent sexual segregation. *Marine Biology* **139**, 1165–1175.
- Stal, J., Pihl, L. and Wennhage, H. (2007) Food utilisation by coastal fish assemblages in rocky and soft bottoms on the Swedish west coast: inference for identification of essential fish habitats. *Estuarine Coastal and Shelf Science* **71**, 593–607.
- Stevens, T. and Connolly, R.M. (2004) Testing the utility of abiotic surrogates for marine habitat mapping at scales

- relevant to management. *Biological Conservation* **119**, 351–362.
- Stoner, A.W. (2003) What constitutes essential nursery habitat for a marine species? A case study of habitat form and function for queen conch *Marine Ecology-Progress Series* **257**, 275–289.
- Stoner, A.W. and Abookire, A.A. (2002) Sediment preferences and size-specific distribution of young-of-the-year Pacific halibut in an Alaska nursery. *Journal of Fish Biology* **61**, 540–559.
- Stoner, A.W. and Titgen, R.H. (2003) Biological structures and bottom type influence habitat choices made by Alaska flatfishes. *Journal of Experimental Marine Biology and Ecology* **292**, 43–59.
- Stoner, A.W., Manderson, J.P. and Pessutti, J.P. (2001) Spatially explicit analysis of estuarine habitat for juvenile winter flounder: combining generalized additive models and geographic information systems. *Marine Ecology-Progress Series* **213**, 253–271.
- Stoner, A.W., Spencer, M.L. and Ryer, C.H. (2007) Flatfish-habitat associations in Alaska nursery grounds: use of continuous video records for multi-scale spatial analysis. *Journal of Sea Research* **57**, 137–150.
- Stoner, A.W., Ryer, C.H., Parker, S.J., Auster, P.J. and Wakefield, W.W. (2008) Evaluating the role of fish behavior in surveys conducted with underwater vehicles. *Canadian Journal of Fisheries and Aquatic Sciences* **65**, 1230–1243.
- Sullivan, M.C., Cowen, R.K., Able, K.W. and Fahay, M.P. (2006) Applying the basin model: assessing habitat suitability of young-of-the-year demersal fishes on the New York Bight continental shelf. *Continental Shelf Research* **26**, 1551–1570.
- Syms, C. (1995) Multiscale analysis of habitat association in a guild of Blennoid fishes. *Marine Ecology-Progress Series* **125**, 31–43.
- Thrush, S.F., Schultz, D., Hewitt, J.E. and Talley, D. (2002) Habitat structure in soft-sediment environments and abundance of juvenile snapper *Pagrus auratus*. *Marine Ecology-Progress Series* **245**, 273–280.
- Valavanis, V.D., Pierce, G.J., Zuur, A.F. et al. (2008) Modelling of essential fish habitat based on remote sensing, spatial analysis and GIS. *Hydrobiologia* **612**, 5–20.
- Van der Kooij, J., Kupschus, S. and Scott, B.E. (2011) Delineating the habitat of demersal fish assemblages with acoustic seabed technologies. *ICES Journal of Marine Science* **68**, 1973–1985.
- Vaz, S., Martin, C., Eastwood, P. et al. (2008) Modelling species distributions using regression quantiles. *Journal of Applied Ecology* **45**, 204.
- Vinagre, C., Fonseca, V., Cabral, H. and Costa, M.J. (2006) Habitat suitability index models for the juvenile soles, *Solea solea* and *Solea senegalensis*, in the Tagus estuary: defining variables for species management. *Fisheries Research* **83**, 140–149.
- Vitousek, P.M. (1997) Human domination of earth's ecosystems. *Science* **277**, 494–499.
- Walsh, S.J., Simpson, M. and Morgan, M.J. (2004) Continental shelf nurseries and recruitment variability in American plaice and yellowtail flounder on the Grand Bank: insights into stock resiliency. *Journal of Sea Research* **51**, 271–286.
- Ward, A., Webster, M. and Hart, P. (2006) Intraspecific food competition in fishes. *Fish and Fisheries* **7**, 231–261.
- Wiens, J.A. (1989) Spatial scaling in ecology. *Functional Ecology* **3**, 385–397.
- Wilhelmsson, D., Malm, T. and Ohman, M.C. (2006) The influence of offshore windpower on demersal fish. *ICES Journal of Marine Science* **63**, 775–784.
- Williams, A. and Bax, N.J. (2001) Delineating fish-habitat associations for spatially based management: an example from the south-eastern Australian continental shelf. *Marine and Freshwater Research* **52**, 513–536.
- Williams, A., Bax, N.J., Kloser, R.J., Althaus, F., Barker, B. and Keith, G. (2009) Australia's deep-water reserve network: implications of false homogeneity for classifying abiotic surrogates of biodiversity. *ICES Journal of Marine Science* **66**, 214–224.
- Wright, D.J. and Heyman, W.D. (2008) Introduction to the special issue: marine and coastal GIS for geomorphology, habitat mapping, and marine reserves. *Marine Geodesy* **31**, 223–230.