

Enterisol[®] Ileitis – Product Characteristics

Description

Enterisol[®] Ileitis is a live attenuated *Lawsonia intracellularis* vaccine for oral drench or drinking water administration

PHARMACEUTICAL FORM

Lyophilisate and diluent for oral suspension

Indications for use

For active immunisation of pigs from three weeks of age and older to reduce intestinal lesions caused by *Lawsonia intracellularis* infection and to reduce growth variability and loss of weight gain associated with the disease.

Benefits

- provides active immunity against the damages caused by *Lawsonia intracellularis* infection.
- significantly improves weight gain
- significantly reduces body weight variability
- protects against diarrhoea and mortality associated with Ileitis
- vaccination can contribute to reduce the total antimicrobial use in pig farms

Advantages

One dose of Enterisol[®] Ileitis offers long lasting immunity instead of need for, continuous use of antibiotics routine.

Dosage and Administration

Administer 1 dose (2 ml) per pig via Drench or in the Drinking water

Contra-indications

None known.

Undesirable effects (frequency and seriousness)

None known.

Special precautions for use

Vaccinate only clinically healthy animals.

Use during pregnancy and lactation

No undesirable effect was observed after administration of the vaccine in breeding and pregnant animals

Interaction with other medicinal products and other forms of interaction

Since the vaccine isolate is a live bacterium, simultaneous use of antimicrobials which are effective against *Lawsonia intracellularis*. should be avoided for a minimum of three days before and after vaccination.

Overdose (symptoms, emergency procedures, antidotes)

No adverse reactions have been observed following administration of ten times the recommended dose.

Special warning for each target species

The vaccine has not been tested in breeding boars; therefore do not vaccinate breeding boars.

Do not vaccinate animals which are receiving treatment with antimicrobials effective against *Lawsonia intracellularis*. Such antimicrobials should be withheld for a minimum of three days before and three days after the day of vaccination.

Withdrawal period(s)

Zero days.

Special precautions for storage

Store and transport between +2°C and +8°C.

Presentations

- 10 dose: (amber glass vial of 20 ml and one polyethylene vial containing 20 ml diluent)
- 50 dose: (amber glass vial of 100 ml and polyethylene vial containing 100 ml diluent)
- 12 x 100 dose: (12 amber glass vials of 100 ml and 12 polyethylene vials containing 200 ml diluent)